

Homework 2
Due 2 May 2018

Problem 1 Incoming solar radiative flux

Given a solar surface temperature of approximately 5800 K, solar radius 7.0×10^5 km, and mean solar distance of 150×10^6 km, calculate the global-mean incoming solar radiative flux at the top of the Earth's atmosphere (TOA).

Problem 2 Radiative equilibrium

Approximate the atmosphere as n black-body layers in radiative equilibrium with each other. (The black-body approximation is going to be increasingly invalid as n increases, but let's see what happens anyway.) The atmosphere is also in radiative equilibrium with a planetary surface of albedo α and TOA solar radiation $S_0/4$. Show that the surface temperature T_s is related to the emission temperature T_e by the relationship

$$T_s = \sqrt[4]{n+1} T_e \quad (1)$$

Is radiative equilibrium a good model for the Earth's atmosphere?

Problem 3 Atmospheric energy budget

In our climate model with a one-layer atmosphere, does the atmosphere experience a net gain or net loss of energy by radiation? What about the real atmosphere?

Problem 4 Stefan–Boltzmann law (optional, for people who like integrals)

Planck's law gives the spectral irradiance from a black body as a function of temperature:

$$B_\lambda(T) d\lambda = \frac{2hc^2}{\lambda^5} \frac{1}{\exp(hc/\lambda k_B T) - 1} d\lambda \quad (2)$$

Consider an infinite plane black body representing the planetary surface, a layer of the atmosphere, or a layer of cloud.

(a) Integrate (2) over a hemisphere to derive the Stefan–Boltzmann law,

$$R = \int_0^\infty d\lambda \int_0^{2\pi} d\phi \int_0^{\pi/2} B_\lambda(T) \cos \theta \sin \theta d\theta = \sigma T^4 \quad (3)$$

- (b) Express σ (the Stefan–Boltzmann constant) in terms of the fundamental constants k_B , c , and h .
- (c) Based on equation (4), what should the size ratio between the two black body curves on p. 5 of the Lecture 2 slides be?

Note 1: You may find it helpful to transform to frequency space.

Note 2: The following integral may be of use:

$$\int_0^{\infty} \frac{x^3}{\exp(x) - 1} dx = \frac{\pi^4}{15} \quad (4)$$

Problem 5 Wien's law (optional, for people who like derivatives)

Show that the spectral radiance $B_\lambda(T)$ peaks at a wavelength proportional to the inverse of the temperature. Find the peak wavelength of a black body at 6000 K and a black body at 255 K.