

Today's Lecture (Lecture 5): General circulation of the atmosphere

Reference

- ▶ Hartmann, *Global Physical Climatology* (1994), Ch. 2, 3, 6
- ▶ Peixoto and Oort, Ch. 4, 6, 7

2.3 – General circulation of the atmosphere

- ▶ Atmospheric transport in response to radiative imbalance
- ▶ Mean meridional circulation and eddy circulation


2.3 – General circulation of the atmosphere

- ▶ Atmospheric transport in response to radiative imbalance
- ▶ Mean meridional circulation and eddy circulation
- ▶ Energy cycle
- ▶ Entropy cycle

2.3 – General circulation of the atmosphere

- ▶ Atmospheric transport in response to radiative imbalance
- ▶ Mean meridional circulation and eddy circulation
- ▶ Energy cycle
- ▶ Entropy cycle
- ▶ Cycles of momentum, angular momentum
- ▶ Hydrological cycle

Radiative budgets for the atmosphere and at TOA



Radiative energy balance of the atmosphere (sign convention: downwelling positive) is


$$R_a = S_{\text{TOA}} - S_s + R_{\text{TOA}} - R_s = (340 - 100) - 160 + (-239) - (342 - 398) \text{ W m}^{-2} = \mathcal{O}(-100 \text{ W m}^{-2}), \quad (2.85)$$

balanced by fluxes of sensible and latent heat into the atmosphere

Zonal-mean radiative budget at TOA


TOA radiative budget...

- ▶ The top-of-atmosphere (TOA) radiative balance measures how much energy enters or leaves the climate system
- ▶ In the tropics, the net energy flux is positive (into the climate system)
- ▶ In the extratropics, the net energy flux is negative (out of the climate system)




... requires meridional energy transport

- ▶ To maintain a steady state (e.g., constant long-term zonal-mean temperature), energy transport from the tropics into the extratropics is required
- ▶ The *meridional divergence* of the energy transport balances the radiative energy flux
- ▶ Contributions to transport from ocean and atmosphere



Atmospheric energy budget requires atmospheric transport




- ▶ As we saw in the previous section, the net radiative energy balance of the atmosphere is $R_a = \mathcal{O}(-100 \text{ W m}^{-2})$; the balance is fairly constant in latitude
- ▶ The radiative energy loss is balanced by latent (LP) and sensible (SH) heat flux from land and ocean; but these are strong functions of latitude
- ▶ Meridional advective atmospheric energy flux is required to provide local energy balance:

$$R_a + F_{LH} + F_{SH} = \Delta F_a \quad (2.86)$$

- ▶ The advective energy flux is the meridional divergence of the meridional heat transport (sign convention: northward positive):

$$\frac{dN}{d\phi} = \int_0^{2\pi} d\lambda R_E^2 \Delta F_a(\phi) \cos \phi = 2\pi R_E^2 \Delta F_a(\phi) \cos \phi \quad (2.87)$$

Atmospheric energy budget requires atmospheric transport



- ▶ As we saw in the previous section, the net radiative energy balance of the atmosphere is $R_a = \mathcal{O}(-100 \text{ W m}^{-2})$; the balance is fairly constant in latitude
- ▶ The radiative energy loss is balanced by latent (LP) and sensible (SH) heat flux from land and ocean; but these are strong functions of latitude
- ▶ Meridional advective atmospheric energy flux is required to provide local energy balance:

$$R_a + F_{LH} + F_{SH} = \Delta F_a \quad (2.86)$$

- ▶ The advective energy flux is the meridional divergence of the meridional heat transport (sign convention: northward positive):

$$\frac{dN}{d\phi} = \int_0^{2\pi} d\lambda R_E^2 \Delta F_a(\phi) \cos \phi = 2\pi R_E^2 \Delta F_a(\phi) \cos \phi \quad (2.87)$$

Averaging operators

Temporal mean

$$\bar{A} = \bar{A}(\lambda, \phi, p) = \frac{1}{\tau} \int_{-\tau/2}^{\tau/2} A(\lambda, \phi, p, t) dt \quad (2.88)$$

and the *zonal mean*

$$[A] = [A](\phi, p, t) = \frac{1}{2\pi} \int_0^{2\pi} A(\lambda, \phi, p, t) d\lambda \quad (2.89)$$

The instantaneous value of A is given by

$$A = \bar{A} + A' \quad (2.90)$$

where A' is called the *fluctuating* component of A . Likewise

$$A = [A] + A^* \quad (2.91)$$

where A^* is the departure from the zonal mean.

Decomposition of a field into time-average and fluctuating, zonally symmetric and zonally asymmetric components:

$$A = [\bar{A}] + [A'] + \bar{A}^* + A'^* \quad (2.92)$$

Decomposition of the flow

Products of fields contain covariance terms (where fluctuations do not average to zero)

$$\overline{AB} = \bar{A}\bar{B} + \overline{A'B'} \quad (2.93)$$

$$[AB] = [A][B] + [A^*B^*] \quad (2.94)$$


$$[\overline{AB}] = [\bar{A}] [\bar{B}] + [\bar{A}^*\bar{B}^*] + [\overline{A'B'}] \quad (2.95)$$

The terms in (2.95) are the **mean circulation**, **stationary eddies**, and **transient eddies**. To take a concrete example, the decomposition of northward flux of sensible heat is

$$c_p [\overline{vT}] = c_p [\bar{v}] [\bar{T}] + c_p [\bar{v}^*\bar{T}^*] + c_p [\overline{v'T'}] \quad (2.96)$$

This week's homework will analyze the relative importance of each contribution as a function of latitude.

Example of meridional heat transport by a transient eddy



$\langle v \rangle [T] \ll [v^* T^*]$ in the eddy covering northern Europe

Mean circulation – streamfunction

The zonal-mean continuity equation (zonal flow is integrated out) is

$$\frac{1}{R_E \cos \phi} \frac{\partial}{\partial \phi} ([\bar{v}] \cos \phi) + \frac{\partial [\bar{\omega}]}{\partial p} = 0 \quad (2.97)$$

For a nondivergent flow, velocity components can be written with the aid of a streamfunction:

$$[\bar{v}] = \frac{g}{2\pi R_E \cos \phi} \frac{\partial \Psi_M}{\partial p} \quad (2.98)$$


$$[\bar{\omega}] = \frac{-g}{2\pi R_E^2 \cos \phi} \frac{\partial \Psi_M}{\partial \phi} \quad (2.99)$$

(2.98) and (2.99) satisfy (2.97); normalization, including the minus sign, is convention – but the relative minus sign is required. To calculate Ψ_M , first impose boundary condition $\Psi_M = 0$ at TOA, then integrate (2.98):

$$\Psi_M = \frac{2\pi R_E \cos \phi}{g} \int_0^p [\bar{v}] dp' \quad (2.100)$$

The normalization is chosen to give units of kg s^{-1} (*mass streamfunction*); the $\cos \phi$ factor is required to ensure constant Ψ_M for constant meridional flow. Mass transport is tangent to contours of the streamfunction. Mass flow between two contours is equal to $\Delta \Psi_M$.


Mean meridional circulation



- ▶ Hadley cell with rising branch in the ITCZ and descending in the subtropics
- ▶ Transport is from winter hemisphere to summer hemisphere at the surface, summer hemisphere to winter hemisphere at altitude → transport of potential energy, latent heat, sensible heat
- ▶ Mass transport by mean circulation is small outside the Hadley cell
- ▶ This is where (temporal and zonal) fluctuations in the circulation are important – eddy transport

Figure: Hartmann (1994); shaded: $\psi_M < 0$; units: $10^{10} \text{ kg s}^{-1}$

Meridional energy transport



- ▶ Recall static energy (2.72): sum of potential energy (PE), sensible heat (SH) and latent heat (LH)

$$h = gz + c_p T + L_v q \quad (2.101)$$

The divergence of poleward transport of these energy terms balances the atmospheric energy budget.

- ▶ Mean transport dominates in the Hadley cell – but note large terms of opposite signs
- ▶ Eddy transport, especially in winter (large temperature gradient), dominates in midlatitudes

