
Quantum Field Theory of Many-Particle Systems - Problem Set 5

Summer Semester 2017

Due: The problem set will be discussed in the tutorial on **Wednesday, 10.05.2017, 13:30**

Internet: The problem sets can be downloaded from
http://home.uni-leipzig.de/stp/QFT_of_MPS_SS17.html

9. Gaussian integrals

1+2+2+1 Punkte

In this problem we will consider different generalisations of the standard Gaussian integral.

a) To begin with show that

$$\int_{-\infty}^{\infty} dx e^{-\frac{1}{2}ax^2} = \sqrt{\frac{2\pi}{a}},$$

where $\text{Re } a > 0$.

b) Generalise the above integral by adding a so-called source term in the exponent. That is calculate

$$\int_{-\infty}^{\infty} dx e^{-\frac{1}{2}ax^2 + Jx}.$$

c) Now let \mathbf{x} and \mathbf{J} be N -dimensional real vectors, and let A be an $N \times N$ matrix. Show that the multi-dimensional analogues of the integrals in parts a and b become

$$\int d\mathbf{x} e^{-\frac{1}{2}\mathbf{x}^T A \mathbf{x}} = (2\pi)^{N/2} \det A^{-1/2},$$

and

$$\int d\mathbf{x} e^{-\frac{1}{2}\mathbf{x}^T A \mathbf{x} + \mathbf{J}^T \cdot \mathbf{x}} = (2\pi)^{N/2} \det A^{-1/2} e^{\frac{1}{2}\mathbf{J}^T A^{-1} \mathbf{J}}.$$

d) What are the analogous expressions if \mathbf{x} and \mathbf{J} are complex vectors and A is a complex matrix?

10. Wick's theorem

2+2+2 Punkte

In this problem we will derive a version of Wick's theorem which plays an important role in quantum field theories. To do this let \mathbf{x} be an N -dimensional real vector and let A be an $N \times N$ matrix. Furthermore define the "expectation value"

$$\langle \dots \rangle = \frac{\int d\mathbf{x} (\dots) e^{-\frac{1}{2}\mathbf{x}^T A \mathbf{x}}}{\int d\mathbf{x} e^{-\frac{1}{2}\mathbf{x}^T A \mathbf{x}}}.$$

a) Using the results from the previous problem, or otherwise, show that

$$\langle x_i x_j \rangle = A_{ij}^{-1},$$

where x_i is the i 'th element of \mathbf{x} and A_{ij}^{-1} is the ij 'th element of the matrix A^{-1} .

b) Likewise show that

$$\langle x_i x_j x_n x_m \rangle = A_{ij}^{-1} A_{nm}^{-1} + A_{in}^{-1} A_{jm}^{-1} + A_{im}^{-1} A_{jn}^{-1}.$$

c) Generalise the above to the product of $2n$ elements of \mathbf{x} .

11. Equation of motion for the Green function of an interacting system

3+5 Punkte

(a) Define the time-ordering operator T for a fermionic and a bosonic system.

(b) Consider the following Hamiltonian in second quantisation,

$$H - \mu N = \int d^3r \psi^\dagger(\mathbf{r}) \epsilon(\mathbf{r}) \psi(\mathbf{r}) + \frac{1}{2} \int d^3r \int d^3r' \psi^\dagger(\mathbf{r}) \psi^\dagger(\mathbf{r}') V(\mathbf{r} - \mathbf{r}') \psi(\mathbf{r}') \psi(\mathbf{r}),$$

where $\epsilon(\mathbf{r}) = -\frac{\hbar^2}{2m} \nabla_{\mathbf{r}}^2 + U(\mathbf{r}) - \mu$ represents the one-particle contribution to the Hamiltonian, and $V(\mathbf{r} - \mathbf{r}')$ is a two-body interaction. Using the Heisenberg equation of motion $\hbar \dot{\psi}(\mathbf{r}, t) = i[H, \psi(\mathbf{r}, t)]$ for the (either fermionic or bosonic) field operator $\psi(\mathbf{r}, t)$, show that the time-ordered Green function $G(\mathbf{r}, t; \mathbf{r}', t') = -i \langle T \psi(\mathbf{r}, t) \psi^\dagger(\mathbf{r}', t') \rangle$ obeys the following equation of motion

$$\left[i\hbar \frac{\partial}{\partial t} - \epsilon(\mathbf{r}) \right] G(\mathbf{r}, t; \mathbf{r}', t') = \hbar \delta(t - t') \delta(\mathbf{r} - \mathbf{r}') - i \int d^3r'' V(\mathbf{r} - \mathbf{r}'') \langle T \psi^\dagger(\mathbf{r}'', t) \psi(\mathbf{r}'', t) \psi(\mathbf{r}, t) \psi^\dagger(\mathbf{r}', t') \rangle.$$

Hint: The following identity for commutators may be useful:

$$[AB, \psi] = A[B, \psi]_{\pm} \mp [A, \psi]_{\pm} B.$$