

Quantum Mechanics 2- Problem Set 7

Wintersemester 2016/2017

Abgabe: The problem set will be discussed in the tutorial on **Monday, 05.12.2016, 13:30**.

19. Continuity equation for the Dirac equation

5 Punkte

Prove the continuity equation

$$\frac{\partial \rho}{\partial t} + \nabla \cdot \mathbf{j} = 0,$$

with

$$\mathbf{j} = \Psi^\dagger \begin{pmatrix} 0 & \sigma \\ \sigma & 0 \end{pmatrix} \Psi,$$

and $\rho = \Psi^\dagger \Psi$ for the Dirac equation.

20. Free particle solutions of the Dirac equation

5 Punkte

Calculate the eigenvalues of the free-particle Dirac equation

$$\begin{pmatrix} m & 0 & p & 0 \\ 0 & m & 0 & -p \\ p & 0 & -m & 0 \\ 0 & -p & 0 & -m \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \\ u_3 \\ u_4 \end{pmatrix} = E \begin{pmatrix} u_1 \\ u_2 \\ u_3 \\ u_4 \end{pmatrix}.$$

21. Klein Tunneling in graphene

1+3+6 Punkte




Figure 1: Left: Schematic drawing of a Dirac electron incident on a potential barrier. Right: Definition of the angles used in the problem.

Consider a Dirac electron with energy E incident on a potential barrier of size V as shown in the figure.

- (a) Why is it sufficient to only require continuity of the wave-function and not its derivative?
- (b) Assume the electron is incident at some angle ϕ in regions I and III and θ in region II, such that $k_x = k \cos \phi$, $k_y = k \sin \phi$ in regions I and III, while $\theta = \arctan(k_y/q_x)$ with $q_x = \sqrt{(V - E)^2/v^2 - k_y^2}$ and $v = k/m$ in region II. Explain why the wave-functions in the different regions can be written as

$$\begin{aligned}\psi_I(x) &= \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ se^{i\phi} \end{pmatrix} e^{i(k_x x + k_y y)} + \frac{r}{\sqrt{2}} \begin{pmatrix} 1 \\ se^{i(\pi-\phi)} \end{pmatrix} e^{i(k_y y - k_x x)}, \\ \psi_{II}(x) &= \frac{a}{\sqrt{2}} \begin{pmatrix} 1 \\ s'e^{i\theta} \end{pmatrix} e^{i(q_x x + k_y y)} + \frac{b}{\sqrt{2}} \begin{pmatrix} 1 \\ s'e^{i(\pi-\theta)} \end{pmatrix} e^{i(k_y y - q_x x)}, \\ \psi_{III}(x) &= \frac{t}{\sqrt{2}} \begin{pmatrix} 1 \\ se^{i\phi} \end{pmatrix} e^{i(k_x x + k_y y)}.\end{aligned}$$

Here $s = \text{sgn}(E)$ and $s' = \text{sgn}(E - V)$. What is the physical significance of r , a , b , and t ?

- (c) Use the continuity of the wave-function to calculate the transmission through the barrier $T(\theta, \phi, Dq_x) = |t|^2$. What do you get for $Dq_x = n\pi$ with n integer? For general values of Dq_x , investigate what happens when $\phi, \theta \rightarrow 0$.