

UNIVERSITÄT LEIPZIG

Climate Dynamics (Summer Semester 2019)
J. Mülmenstädt

Today's Lecture (Lecture 13): Uncertainties due to clouds and aerosols


Reference

IPCC AR5, Chs. 7 and 9


5.3 – Clouds and aerosols

Why discuss clouds aerosols and clouds together?

- ▶ Anthropogenic activity affects clouds through aerosols and GHG (both through the surface temperature and through rapid adjustments)
- ▶ Past: strong aerosol forcing (relative to GHG); inter-model spread dominated by differences in aerosol
- ▶ Future: weak aerosol forcing (relative to GHG); inter-model spread dominated by feedbacks, mainly cloud
- ▶ Inability to constrain climate sensitivity from historical observations if the aerosol ERF is poorly constrained
- ▶ Clouds and aerosols each pose two distinct challenges:
 1. Fundamental understanding of processes
 2. Their representation in large-scale models



Challenges related to clouds



Representation of clouds in climate models


Parameterized **subgridscale** processes:

- ▶ Turbulence
- ▶ Cumulus convection
- ▶ Microphysical processes
- ▶ Radiative transfer
- ▶ Cloud amount (including the vertical overlap between different grid levels)
- ▶ Subgridscale transport of aerosol and chemical species

Many cloud processes are unrealistic in current GCMs → cloud response to climate change remains uncertain

CRE is large compared to feedbacks (and forcings)

- ▶ LW and SW CRE: $\mathcal{O}(10) \text{ W m}^{-2}$
- ▶ Forcings: $\mathcal{O}(1) \text{ W m}^{-2}$
- ▶ Feedbacks: $\mathcal{O}(1) \text{ W m}^{-2} \text{ K}^{-1}$




Figures: IPCC AR5 unless noted

Need for models to evaluate feedbacks

- ▶ Observable climate variations are not necessarily good analogs for GHG climate change
- ▶ Change in TOA flux due to clouds is difficult to isolate


Feedbacks: water vapor + lapse rate



Compensation in intermodel spread of water vapor and lapse rate feedback

- ▶ Saturation water vapor pressure as a function of surface temperature: $7\% \text{ K}^{-1}$ near the surface, up to $17\% \text{ K}^{-1}$ in the upper troposphere
- ▶ Increase with height because of the lapse rate feedback
- ▶ Models with strong lapse rate feedbacks will have high increase in upper tropospheric water vapor, and therefore a strong water vapor feedback
- ▶ Combined lapse rate + water vapor feedback is well constrained; $+0.96$ to $+1.22 \text{ W m}^{-2} \text{ K}^{-1}$

Feedbacks: clouds




Cloud feedbacks:

- ▶ Changes in high-level cloud altitude and amount
- ▶ Effects of hydrological cycle and storm track changes on cloud systems
- ▶ Changes in low-level cloud amount
- ▶ Microphysically induced opacity (optical depth) changes
- ▶ Changes in high-latitude clouds


Some changes occur at the GCM resolved scale, but most involve **subgrid-scale processes** that need to be parameterized

Cloud feedbacks: high-cloud altitude




- ▶ Ascent in tropical deep convection is mass-balanced by compensating subsidence
- ▶ Compensating subsidence is due to equilibrium between radiative cooling and adiabatic compression
- ▶ The subsidence top occurs at the altitude where the water vapor mixing ratio decreases rapidly ($\approx 220 \text{ K}$); the convection top will occur at the same altitude
- ▶ In a warming climate, the water vapor mixing ratio still has the same temperature dependence, so that the radiative cooling still become inefficient at $\approx 220 \text{ K}$
- ▶ The clear-sky emission temperature will increase due to atmospheric warming, but the cloud emission temperature will not, so that the LW CRE becomes stronger
- ▶ Expect $+0.5 \text{ W m}^{-2} \text{K}^{-1}$ (in the tropics); model range is $+0.09$ to $+0.58 \text{ W m}^{-2} \text{K}^{-1}$

Cloud feedbacks: circulation changes



Boundary layer – the cloud-process view



Vertical structure

Boundary layer is well mixed and capped by a ...

Cloud layer which maintains a temperature inversion by cloud-top cooling and is weakly coupled to the ...

Free troposphere by an entrainment layer

Processes

Sensible and latent heat flux at the surface and ...


Radiative cooling at cloud top destabilize the airmass; this results in ...

Convection which mixes the layer vertically and horizontally

Cloud feedbacks: low cloud


radiative

↓ more emissive FT
(more CO₂ or H₂O)




thermodynamic

warmer SST
or drier RH




dynamic

↓ ↓ ↓
less subsidence



inversion strength

FT warms more than SST



Cloud feedbacks: low cloud

Low clouds, especially in the tropics and subtropics, are the largest contributors to the intermodel spread in cloud feedback

Negative feedback mechanisms

In a warmer climate, low clouds might be


- ▶ horizontally more extensive, because changes in the lapse rate also modify the lower-tropospheric stability
- ▶ optically thicker, because adiabatic ascent condenses more liquid
- ▶ vertically more extensive in response to weakening of the tropical overturning circulation

Positive feedback mechanisms


- ▶ Warming-induced increase in moisture inversion strength reduces cloud amount or thickness
- ▶ Energetic constraints prevent the surface evaporation from increasing with warming at a rate sufficient to balance expected changes in dry air entrainment, thereby reducing the supply of moisture to form clouds
- ▶ Increased concentrations of GHGs reduce the radiative cooling that drives stratiform cloud layers and thereby the cloud amount

It appears that the positive feedbacks, though less intuitive, are more important; in GCMs, the low-cloud feedback ranges from -0.09 to $+0.63 \text{ W m}^{-2} \text{ K}^{-1}$ (with approximately 80% probability of positive feedback); high-resolution modeling supports the mechanisms above

Radiative forcing: aerosol-radiation and aerosol-cloud interactions



Aerosol–cloud and aerosol–radiation interactions: large uncertainties



Confounding by meteorology

Aerosol depends on air mass history (origin, precipitation, humidity, . . .), **but so do clouds**

Non-monotonic behavior of the adjustments

Magnitude and even sign of the adjustments depends on details of small-scale processes

Uncertain preindustrial state

Unlike for WMGHG, we have no reliable estimates of preindustrial aerosol; biomass burning contributed anthropogenic aerosol even before the Industrial Revolution

Category	Best Estimate	Climate Model and/or Satellite Instrument	Reference
with mixed-phase clouds	-1.55	CAM Oslo	Hoose et al. (2010b)
with mixed-phase clouds	-1.02	ECHAM	Lohmann and Ferrachat (2010)
with mixed-phase clouds	-1.68	GFDL	Salzmann et al. (2010)
with mixed-phase clouds	-0.81	CAM Oslo	Storelvmo et al. (2008b, 2010)
with convective clouds	-1.50	ECHAM	Lohmann (2008)
with convective clouds	-1.38	GISS	Koch et al. (2009a)
with convective clouds	-1.05	PNL-MMF	Wang et al. (2011b)
Satellite-based	-0.85	ECHAM + POLDER	Lohmann and Lesins (2002)
Satellite-based	-0.93	AVHRR	Sekiguchi et al. (2003)
Satellite-based	-0.67	CERES / MODIS	Lebeck et al. (2008)
Satellite-based	-0.45	CERES / MODIS	Quaas et al. (2008)
Satellite-based	-0.95	Model mean + MODIS	Quaas et al. (2009)
Satellite-based	-0.85	MACC + MODIS	Bellouin et al. (2013)

AVHRR = Advanced Very High Resolution Radiometer. MACC = Monitoring Atmospheric Composition and Climate. POLDER = Polarization and Directionality of the Earth's Reflectances. CERES = Clouds and the Earth's Radiant Energy System. MODIS = Moderate Resolution Imaging Spectrometer.